

# CME arrival-time validation of real-time WSA-ENLIL+Cone simulations at the CCMC/SWRC

Alexandra M. Wold[1], M. Leila Mays[2,3], A. Taktakishvili[2,3], L. Jian[4,3], D.Odstrcil[2,5], P. MacNeice[3]

[1] American University, [2] Catholic University of America, [3] NASA Goddard Space Flight Center, [4] University of Maryland College Park, [5] George Mason University

## **METHODOLOGY**


- ➤ Validation results of the WSA-ENLIL+Cone model running in real-time at the CCMC/SWRC and archived in DONKI.
- ➤ Compare model predicted CME arrival-times to in-situ ICME shock observations near Earth, STEREO-A & B, March 2010 present (simulations of 2,700 CMEs).
- > Report hit, miss, false alarm, and correct rejection statistics
  - ➤ For hits, compute bias and average absolute arrival time error and dependence of errors on CME input parameters
- ➤ Determine **impact of multi-spacecraft observations** on the CME parameters used to initialize the model
  - ➤ before and after the STEREO B communication loss (since September 2014) & STEREO-A side-lobe operations (August 2014-December 2015)

#### **VALIDATION PROCESS**

- ➤ The quality of model operation is evaluated by comparing the model output to the observed CME arrival time.
  - Referencing ICME catalogues
     (Lan Jian, Teresa Nieves,
     Richardson & Cane, ISEST,
     HELCATS)
  - ➤ DONKI database
  - ➤ Analysis of in situ data (ACE, WIND, STEREO)
- ➤ Complications:
  - ➤ Observed arrival is weak
  - ➤ Hybrid SIR and CME event
  - ➤ CME arrival with uncertain source.











#### **CONTINGENCY TABLE & SKILL SCORES**


- ➤ Success Ratio: H / (H+FA)
  - ➤ fraction of predicted arrivals that were correct; perfect score = 1
- ➤ False Alarm Ratio: FA / (H+FA)
  - ➤ fraction of predicted arrivals that were incorrect; perfect score = 0
- ➤ Accuracy: (H+CR) / Total
  - ➤ fraction of correct forecasts; perfect score = 1
- ➤ Bias Score: (H+FA) / (H+M)
  - ➤ ratio of predicted arrivals to observed arrivals; under forecast <1; perfect score = 1; over forecast >1
- ➤ Probability of Detection (hit rate): H / (H+M)
  - ➤ fraction of observed events that were predicted; perfect score = 1

|                      | Observed Arrival | No Observed Arrival    |
|----------------------|------------------|------------------------|
| Predicted Arrival    | Hit (H)          | False Alarm (FA)       |
| No Predicted Arrival | Miss (M)         | Correct Rejection (CR) |

## HITS, FALSE ALARMS, MISSES & CORRECT REJECTIONS


|                           | Earth | STEREO-A | STEREO-B | All  |
|---------------------------|-------|----------|----------|------|
| Hits                      | 128   | 105      | 58       | 291  |
| False Alarms              | 112   | 85       | 53       | 250  |
| Misses                    | 109   | >108     | >71      | >288 |
| <b>Correct Rejections</b> | 1114  | 1195     | 848      | 3157 |

#### Success and False Alarm Ratios




### CME ARRIVAL TIME PREDICTION ERRORS


#### Average Absolute Error at Earth, STEREO A & B




## CME ARRIVAL TIME ERROR DISTRIBUTION




#### CME arrival time error compared to input speed



Input CME radial speed (km/s)
CME arrival time error compared to input speed




# Kp Range Prediction Skill Scores Is the observed Kp<sub>max</sub> within the predicted Kp range?




#### Kp skill scores varying with threshold

Using threshold varying contingency tables for observed and predicted  $Kp_{max}$ 



#### Error in predicting $Kp_{max}$ varied by threshold



## **CONCLUSIONS**

- ➤ Simulations of 2,700 CMEs (March 2010 present), 291 observed arrivals:
  - ➤ CME arrival time average absolute error at all locations = 10 hours
- ➤ Preliminary validation of the period post-September 2014 (without STEREO B and reduced STEREO A coverage) shows a reduction in skill of **2.3 hours** (36 observed arrivals)
  - multi-view coronagraph observations have a quantitative impact on CME arrival time forecast accuracy (support for L5 need)
- ➤ Sources of CME arrival time error:
  - ➤ input CME parameters
  - ambient solar wind prediction (and magnetogram limitations/ uncertainties)
  - ➤ input model ambient parameters
  - > model assumptions CME has no flux-rope field

#### **FUTURE WORK**

- ➤ Introduce quality factors for observed arrivals and identifying candidate CMEs
- ➤ Look at other locations (MESSENGER, Mars)
  - ➤ Winslow et al. (2015) MESSENGER ICME catalogue